

Hadron physics: a quark-model analysis

A. Valcarce*, J. Vijande[†], P. González** and H. Garcilazo[‡]

**Dpto. de Física Fundamental, Universidad de Salamanca, Spain*

[†]Dpto. de Física Teórica, Universidad de Valencia (UV) and IFIC (UV-CSIC), Spain

***Dpto. de Física Teórica, Universidad de Valencia, Spain*

[‡]Escuela Superior de Física y Matemáticas, Instituto Politécnico Nacional, Mexico

Abstract. We discuss recent results on heavy and light baryon spectroscopy.

Keywords: baryons, quark-model

PACS: 12.39.Jh, 12.39.Pn, 14.20.-c

INTRODUCTION

Hadron spectroscopy has undergone a great renaissance in recent years [1]. The new findings include: low-lying excitations of D and B mesons, long-awaited missing states and new states near 4 GeV/c² in the charmonium spectrum, charmed and bottom baryons, and evidence for doubly charmed baryons. The light hadron sector remains also restless reporting new scalar mesons or showing a deep theoretical interest in the high energy part of both the meson and baryon spectra. In this talk we center our attention in the heavy baryon spectroscopy as well as some anomalous states present in the light baryon spectra.

HEAVY BARYONS

Heavy baryons containing a single heavy quark are particularly interesting. The light degrees of freedom (quarks and gluons) circle around the nearly static heavy quark. Such a system behaves as the QCD analogue of the familiar hydrogen bounded by the electromagnetic interaction. When the heavy quark mass $m_Q \rightarrow \infty$, the angular momentum of the light degrees of freedom is a good quantum number. Thus, heavy quark baryons belong to either SU(3) antisymmetric $\mathbf{\bar{3}}_F$ or symmetric $\mathbf{6}_F$ representations. The spin of the light diquark is 0 for $\mathbf{\bar{3}}_F$, while it is 1 for $\mathbf{6}_F$. Thus, the spin of the ground state baryons is 1/2 for $\mathbf{\bar{3}}_F$, representing the Λ_b and Ξ_b baryons, while it can be both 1/2 or 3/2 for $\mathbf{6}_F$, allocating Σ_b , Σ_b^* , Ξ_b' , Ξ_b^* , Ω_b and Ω_b^* , where the star indicates spin 3/2 states. Therefore heavy hadrons form doublets. For example, Σ_b and Σ_b^* will be degenerate in the heavy quark limit. Their mass splitting is caused by the chromomagnetic interaction at the order $1/m_Q$. These effects can be, for example, taken into account systematically in the framework of heavy quark effective field theory. The mass difference between states belonging to the $\mathbf{\bar{3}}_F$ and $\mathbf{6}_F$ representations do also contain the dynamics of the light diquark subsystem, hard to accommodate in any heavy quark mass expansion. Therefore, exact solutions of the three-body problem for

TABLE 1. Masses, in MeV, of charmed and bottom baryons.

State	J^P	Charm					Bottom				
		CQC	Exp.	[3]	[4]	[5]	CQC	Exp.	[3]	[4]	[5]
Λ_i	$1/2^+$	2285	2286	2285	2268	2297	5624	5624	5638	5612	5622
	$1/2^+$	2785	2765	2865	2791	2772	6106		6188	6107	6086
	$1/2^-$	2627	2595	2635	2625	2598	5947		5978	5939	5930
	$1/2^-$	2880	2880	2885	2816	3017	6245		6268	6180	6328
	$3/2^+$	3061		2930	2887	2874	6388		6248	6181	6189
	$3/2^+$	3308		3160	3073	3262	6637		6488	6401	6540
	$5/2^+$	2888	2880	2930	2887	2883					
Σ_i	$1/2^+$	2435	2454	2455	2455	2439	5807	5808	5845	5833	5805
	$1/2^+$	2904		3025	2958	2864	6247		6370	6294	6202
	$1/2^-$	2772	2765	2805	2748	2795	6103		6155	6099	6108
	$1/2^-$	2893		2885		3176	6241		6245		6401
	$3/2^+$	2502	2518	2535	2519	2518	5829	5829	5875	5858	5834
	$3/2^+$	2944	2940	3065	2995	2912	6260		6385	6308	6222
	$3/2^-$	2772	2800	2805	2763	2761					
Ξ_i	$1/2^+$	2471	2471	2467	2492	2481	5801	5793	5806	5844	5812
	$1/2^+$	3137	3123	2992		2923	6258		6306		6264
	$1/2'^+$	2574	2578	2567	2592	2578	5939		5941	5958	5937
	$1/2'^+$	3212		3087		2984	6360		6416		6327
	$1/2^-$	2799	2792	2792	2763	2801	6109		6116	6108	6119
	$1/2^-$	2902		2897	2859	2928	6223		6236	6192	6238
	$1/2^-$	3004	2980	2993		3186					
	$3/2^+$	2642	2646	2647	2650	2654	5961		5971	5982	5963
	$3/2^+$	3071	3076	3057	2984	3030	6373		6356	6294	6341
	$5/2^+$	3049	3055	3057		3042					
	$5/2'^+$	3132	3123	3167		3123					
Ω_i	$1/2^+$	2699	2698	2675	2718	2698	6056		6034	6081	6065
	$1/2^+$	3159		3195	3152	3065	6479		6504	6472	6440
	$1/2^-$	3035		3005	2977	3020	6340		6319	6301	6352
	$1/2^-$	3125		3075		3371	6458		6414		6624
	$3/2^+$	2767	2768	2750	2776	2768	6079		6069	6102	6088
	$3/2^+$	3202		3235	3190	3119	6493		6519	6478	6518

heavy hadrons are theoretically desirable because they will serve to test the reliability of approximate techniques: heavy quark mass expansions, variational calculations, or quark-diquark approximations.

We have solved the Schrödinger equation by the Faddeev method in momentum space with the constituent quark model (CQC) of Ref. [2]. The results are shown in Table 1 compared to experiment and other theoretical approaches. All known experimental data are nicely described. Such an agreement and the exact method used to solve the three-body problem make our predictions also valuable as a guideline to experimentalists.

As compared to other results in the literature we see an overall agreement for the low-lying states both with the quark-diquark approximation of Ref. [5] and the variational calculation in a harmonic oscillator basis of Ref. [4]. It is worth noticing that the relativistic quark-diquark approximation and the harmonic oscillator variational method predict a lower $3/2^+$ excited state for the Λ_b baryon. Such result can be easily under-

TABLE 2. Masses, in MeV, of different bottom baryons with two-light quarks with (Full) and without ($V_\pi = 0$) the contribution of the one-pion exchange potential.

State	Full	$V_\pi = 0$	ΔE
$\Sigma_b(1/2^+)$	5807	5822	-15
$\Sigma_b(3/2^+)$	5829	5844	-15
$\Lambda_b(1/2^+)$	5624	5819	-195
$\Lambda_b(3/2^+)$	6388	6387	+ 1

stood by looking at Table 2, where it is made manifest the influence of the pseudoscalar interaction between the light quarks on the $\Lambda_b(1/2^+)$ ground state, diminishing its mass by 200 MeV. If this attraction were not present for the $\Lambda_b(1/2^+)$, the $\Lambda_b(3/2^+)$ it would be lower in mass as reported in Refs. [4, 5] (a similar effect will be observed in the charmed baryon spectra). Thus, the measurement and identification of the $\Lambda_b(3/2^+)$ is a relevant feature that will help to clarify the nature of the interaction between the light quarks in heavy baryon spectroscopy, determining the need of pseudoscalar forces consequence of the spontaneous chiral symmetry breaking in the light flavor sector.

In the case of charmed baryons, there are some excited states that it is not even known if they are excitations of the Λ_c or Σ_c . Besides, a number of new Ξ_c and Ξ'_c states have been also discovered recently [2]. As can be seen all known experimental states fit nicely into the description of our model not leaving too many possibilities open for the assigned quantum numbers as we resume in Table 3.

Finally, we can make parameter free predictions for ground states as well as for spin, orbital and radial excitation of doubly charmed and bottom baryons. Our results are shown in Table 4. For doubly charmed baryons, the ground state is found to be at 3579 MeV, far below the result of Ref. [4] and in perfect agreement with lattice nonrelativistic QCD [6], but still a little bit higher than the non-confirmed SELEX result, 3519 MeV [7].

TABLE 3. Possible model states and spin-parity assignments for recently discovered charmed baryons. The 'star' indicates radial excitations.

Experimental resonance (MeV)	Model states
Λ_c or Σ_c	
2765	$\Sigma_c(1/2^-)$ or $\Lambda_c(1/2^+)^*$
2880	$\Lambda_c(1/2^-)^*$ or $\Lambda_c(5/2^+)$
2940	$\Sigma_c(3/2^+)^*$
2800	$\Sigma_c(3/2^-)$
Ξ_c or Ξ'_c	
3055	$\Xi_c(5/2^+)$
3123	$\Xi_c(1/2^+)^*$ or $\Xi'_c(5/2^+)$
2980	$\Xi_c(1/2^-)^*$
3076	$\Xi_c(3/2^+)^*$

TABLE 4. Ground state and excitation energies, ΔE , of doubly charmed and bottom baryons. The 'star' indicates radial excitations. Masses are in MeV.

State	J^P	CQC	[8]	[3]	[6]	[4]
Ξ_{bb}	$1/2^+$	10189	10340	10194		10340
	$3/2^+$	29	30	41	20	27
	$3/2^{+*}$	312		386		238
	ΔE $1/2^{+*}$	293		355		236
	$1/2^-$	217		262		153
	$1/2^{-*}$	423		462		370
Ω_{bb}	$1/2^+$	10293	10370	10267		10454
	$3/2^+$	28	30	38	19	32
	$3/2^{+*}$	329		383		267
	ΔE $1/2^{+*}$	311		359		239
	$1/2^-$	226		265		162
	$1/2^{-*}$	390		410		309
Ξ_{cc}	$1/2^+$	3579	3660	3607	3588	3676
	$3/2^+$	77	80	93	70	77
	$3/2^{+*}$	446		486		366
	ΔE $1/2^{+*}$	397		435		353
	$1/2^-$	301		314		234
	$1/2^{-*}$	439		472		398
Ω_{cc}	$1/2^+$	3697	3740	3710	3698	3815
	$3/2^+$	72	40	83	63	61
	$3/2^{+*}$	463		498		373
	ΔE $1/2^{+*}$	415		445		365
	$1/2^-$	312		317		231
	$1/2^{-*}$	404		410		320

It is therefore a challenge for experimentalists to confirm or to find the ground state of doubly charmed and bottom baryons.

The combined study of Qqq and QQq systems, where Q stands for a heavy c or b quark and q for a light u , d , or s quark, will also provide some hints to learn about the basic dynamics governing the interaction between light quarks. The interaction between pairs of quarks containing a heavy quark Q is driven by the perturbative one-gluon exchange. For the Qqq system the mass difference between members of the $\mathbf{6_F}$ SU(3) representation comes determined only by the perturbative one-gluon exchange, whether between members of the $\mathbf{6_F}$ and $\mathbf{3_F}$ representations it presents contributions from the one-gluon exchange and also possible pseudoscalar exchanges. If the latter mass difference is attributed only to the one-gluon exchange (this would be the case of models based only on the perturbative one-gluon exchange), it will be strengthened as compared to models considering pseudoscalar potentials at the level of quarks, where a weaker one-gluon exchange will play the role. When moving to the QQq systems

only one-gluon exchange interactions between the quarks will survive, with the strength determined in the $Q\bar{q}q$ sector, where we have experimental data. This will give rise to larger masses for the ground states, due to the more attractive one-gluon exchange potential in the $Q\bar{q}q$ sector, what requires larger constituent quark masses to reproduce the experimental data. This could be the reason for the larger masses of ground state doubly heavy baryons obtained with gluon-based interacting potentials [4, 9].

LIGHT BARYONS

In the Particle Data Group (PDG) book [10] the light-quark (u and d) baryon spectrum is composed of forty resonances rated from one (*) to four (****) stars. The PDG average-mass region below 1950 MeV contains mostly four-star (well established) resonances, fourteen out of twenty three, the same being true for the Λ strange sector, eight out of eleven. This makes this mass region the most suitable for testing any spectroscopic quark model. From the pioneering Isgur and Karl's non-relativistic quark model in the late 70's [11] more refined spectroscopic quark models for baryons, based on two-body interactions, have been developed [12]. We will refer to them as two-body quark models and we shall denote them generically as $3q^{2b}$. As an overall result the masses of the fourteen four-star resonances, most times with the exception of $N_{P_{11}}(1440)$, are rather well predicted (≤ 100 MeV difference with the PDG average value) by these models. Regarding the five three-star (likely to certain existence) resonances, the situation is much less favorable since the masses of two of them, $\Delta_{P_{33}}(1600)$ and $\Delta_{D_{35}}(1930)$, are generally overpredicted, up to 250 MeV above the PDG average value. Let us note that a similar discrepancy is observed for $\Delta_{S_{31}}(1900)(**)$ and $\Delta_{D_{33}}(1940)(*)$ (≥ 100 MeV difference with the PDG average value) which can be related to $\Delta_{D_{35}}(1930)$ as we shall show, and for $\Delta_{P_{31}}(1750)(*)$ (up to 200 MeV above the PDG average). In the strange Λ sector an outstanding overpredicted (by 80 – 150 MeV) state is the $\Lambda_{S_{01}}(1405)(****)$. Henceforth we shall call anomalies these significantly overpredicted mass resonances.

We carry out a general analysis of the anomalies: we identify them and we propose a plausible physical mechanism to give correctly account of their masses. Among the anomalies we find large-energy-step anomalies, that correspond either to radial excitations as the $\Delta_{P_{33}}(1600)$ and the $N_{P_{11}}(1440)$ or quark Pauli blocking induced states as the $\Delta_{D_{35}}(1930)$ and the $\Delta_{P_{31}}(1750)$ [12].

Given the large radial excitation energy and the large mass predicted for quark Pauli blocking induced states, one may wonder about the possibility that $4q1\bar{q}$ components may be energetically competitive, despite the extra quark and antiquark masses. Thus, they could greatly contribute, altogether with $3q$ components, to the formation of the bound structures. In order to examine this possibility at a phenomenological level we look for $4q1\bar{q}$ components in the form of inelastic meson-baryon channels in relative S wave (the lowest energy partial wave) with adequate quantum numbers to couple to the anomalies and with thresholds close above their PDG masses. We shall name these components meson-baryon threshold channels or mB channels.

Certainly meson-baryon channel coupling effects may be at work for other resonances not involving either large energy excitation steps or a large mass induced by quark Pauli blocking. The most prominent examples are the $\Lambda_{S_{01}}(1405)$ being mostly interpreted, at

the hadron level, as an S wave $N\bar{K}$ quasi-bound system, and the $\Delta_{F_{35}}(2000)$, a bizarre state since its average mass is obtained from three different data analyses, two of them [13] reporting a mass about $1720 (\pm 60)$ MeV and the other [14] giving a quite different value of 2200 ± 125 MeV. Then by considering two differentiated resonances the $\Delta_{F_{35}}(\sim 1720)$ would be a clear candidate for an anomaly.

To go beyond the qualitative analysis of the anomalies we shall consider a system of one confined channel, the $3q^{2b}$, in interaction with one free-channel, a meson-baryon threshold channel mB , with a hamiltonian matrix:

$$[H] \simeq \begin{pmatrix} M_m + M_B & a \\ a^* & M_{3q^{2b}} \end{pmatrix} \quad (1)$$

where $M_{3q^{2b}}$ stands for the mass of the $3q^{2b}$ state, M_m and M_B for the masses of the meson and baryon respectively and a for a fitting parameter giving account of the interaction.

In order to proceed to calculate the eigenvalues we have to choose a particular $3q^{2b}$ model and establish a criterion for the choice of the mB channel for each anomaly. We shall use as $M_{3q^{2b}}$ the values calculated in Ref. [9]. As mB we shall take for granted the $N\bar{K}$ channel for $\Lambda_{S_{01}}(1405)$ and the σN channel for $N_{P_{11}}(1440)$. For $\Delta_{D_{35}}(1930)$, $\Delta_{D_{33}}(1940)$ and $\Delta_{S_{31}}(1900)$ we shall select $\rho\Delta$ (equivalently we could have preferred the almost degenerate $\omega\Delta$) as suggested by our phenomenological analysis. For the same reason $\pi N_{D_{15}}(1675)$ will be employed for $\Delta_{F_{35}}(\sim 1720)$. For $\Delta_{P_{31}}(1750)$ we shall use $\pi N_{S_{11}}(1650)$ and for $\Delta_{P_{33}}(1600)$ we shall take $\pi N_{P_{13}}(1520)$.

Although the value of $|a|$ might vary depending on the configurations involved in each $(mB) - 3q$ coupling we shall use for the sake of simplicity the same value in all cases. The M results for $|a| = 85$ MeV are numerically detailed in Table 5 where the values for $M_{3q^{2b}}$ and for $(M_m + M_B)$ in the chosen mB channel as well as their probabilities to give M are also displayed. As can be checked the improvement of the description with respect to $3q^{2b}$ is astonishing. All the predicted M masses lye very close to the PDG average masses for the anomalies. In Fig. 1 the M values for $|a| = 85$ MeV are drawn as compared to the experimental mass intervals.

TABLE 5. Predicted masses, M , for the anomalies as compared to experimental data from Ref. [10], Ref. [14] (indicated by the superindex $\dagger$), and Ref. [13] (indicated by the superindex $\dagger\dagger$). Two-body quark-model masses ($3q^{2b}$) are taken from Ref. [9]. Probabilities (Prob.) for meson-baryon and $3q$ components are also shown. All masses are in MeV.

PDG Resonance	mB threshold	Prob.	$3q^{2b}$	Prob.	M	Experiment
$\Delta_{P_{33}}(1600)(***)$	$[\pi N_{P_{13}}(1520)]$ (1660)	81.1%	1795	18.9%	1619	1550–1700
$N_{P_{11}}(1440)(****)$	$[\sigma N]$ (1540)	50.0%	1540	50.0%	1455	1420–1470
$\Delta_{D_{35}}(1930)(***)$	$[\rho\Delta]$ (2002)	83.4%	2155	16.6%	1964	1900–2020
$\Delta_{D_{33}}(1940)(*)$		82.2%	2145	17.8%	1962	1840–2040 †
$\Delta_{S_{31}}(1900)(*)$		81.5%	2140	18.5%	1961	1850–1950
$\Delta_{P_{31}}(1750)(*)$	$[\pi N_{S_{11}}(1650)]$ (1790)	62.8%	1835	37.2%	1725	1710–1780
$\Delta_{F_{35}}(\approx 1720)(N.C.)$	$[\pi N_{D_{15}}(1675)]$ (1815)	74.4%	1910	25.6%	1765	1660–1785 ††
$\Lambda_{S_{01}}(1405)(****)$	$[\bar{K}N]$ (1434)	78.2%	1550	21.8%	1389	1400–1410




FIGURE 1. Predicted masses for the anomalies (dashed lines) as compared to the experimental mass intervals detailed in Table 5 (boxes). N.C. means non-cataloged resonance.

We interpret these results as providing strong quantitative support to our former qualitative description of the anomalies. Regarding their nature a look at the probabilities reveal they are mostly meson-baryon states. Actually a meson-baryon probability greater or equal than 50% can serve as a criterion to identify an anomaly. Nonetheless the coupling to the $3q$ component is essential to lower their masses making them more stable against decay into $m + B$.

It should be emphasized that similar results could be obtained for any other spectroscopic $3q^{2b}$ model through a fine tuning of the value of $|a|$ (note that the small value of $|a|$ as compared to $M_{3q^{2b}}$ and $(M_m + M_B)$ provides an *a posteriori* validation of our method). This comes from the expression of the eigenvalues where it is clear that even for $|a| = 0$ one gets $M = M_m + M_B$ which according to our mB choice is much closer to the PDG mass of the anomaly than $M_{3q^{2b}}$. This means that concerning the mass of the anomalies the coupling of meson-baryon to $3q$ components may play the role of a general *healing mechanism* for spectroscopic models.

SUMMARY

We have studied the heavy baryon spectra by means of the Faddeev method in momentum space. These results should be highly valuable both from the theoretical and experimental points of view. Theoretically, it should be a powerful tool for testing different

approximate methods to solve the three-body problem. Experimentally, the remarkable agreement with known experimental data make our predictions highly valuable as a guideline to experimentalists.

Heavy baryons constitute an extremely interesting problem joining the dynamics of light-light and heavy-light subsystems in an amazing manner. While the mass difference between members of the same SU(3) configuration, either $\mathbf{3}_F$ or $\mathbf{6}_F$, is determined by the perturbative one-gluon exchange, the mass difference between members of different representations comes mainly determined by the dynamics of the light diquark, and should therefore be determined in consistency with the light baryon spectra. There is therefore a remnant effect of pseudoscalar forces in the two-light quark subsystem.

For light baryons we propose that $4q1\bar{q}$ components, in the form of S wave meson-baryon channels which we identify, play an essential role in the description of the anomalies, say baryon resonances very significantly overpredicted by three-quark models based on two-body interactions. As a matter of fact by considering a simplified description of the anomalies as systems composed of a free meson-baryon channel interacting with a three-quark confined component we have shown they could correspond mostly to meson-baryon states but with a non-negligible $3q$ state probability which makes their masses to be below the meson-baryon threshold. The remarkable agreement of our results with data in all cases suggests the implementation of meson-baryon threshold effects as an essential physical mechanism to give account of spectral states poorly described by constituent quark models.

ACKNOWLEDGMENTS

This work has been partially funded by the Spanish MEC and EU FEDER under Contract No. FPA2007-65748, by EIII 506078, by JCyL under Contract No. SA016A17, and by COFAA-IPN (México).

REFERENCES

1. J. L. Rosner, *J. Phys. Conf. Ser.* **69**, 012002 (2007).
2. A. Valcarce, H. Garcilazo, and J. Vijande *Eur. Phys. J. A*, in print (2008) and references therein.
3. B. Silvestre-Brac, *Few-Body Systems* **20**, 1 (1996).
4. W. Roberts and M. Pervin, *arXiv:0711.2492*.
5. D. Ebert, R. N. Faustov, and V. O. Galkin, *Phys. Lett. B* **659**, 612 (2008).
6. N. Mathur, R. Lewis, and R. M. Woloshyn, *Phys. Rev. D* **66**, 014502 (2002).
7. SELEX Collaboration, A. Ocherashvili *et al.*, *Phys. Lett. B* **628**, 18 (2005).
8. R. Roncaglia, D. B. Lichtenberg, and E. Predazzi, *Phys. Rev. D* **52**, 1722 (1995).
9. S. Capstick and N. Isgur, *Phys. Rev. D* **34**, 2809 (1986).
10. W. -M. Yao *et al.*, *J. Phys. G* **33**, 1 (2006).
11. N. Isgur and G. Karl, *Phys. Lett.* **72B**, 109 (1977); *Phys. Rev. D* **19**, 2653 (1979).
12. P. González, J. Vijande, and A. Valcarce, *Phys. Rev. C* **77**, 065213 (2008) and references therein.
13. T. P. Vrana, S. A. Dytman, and T. -S. H. Lee, *Phys. Rep.* **328**, 181 (2000).
14. R. E. Cutkosky, C. P. Forsyth, J. B. Babcock, R. L. Kelly, and R. E. Hendrick, Proceedings of the IV International Conference on Baryon Resonances (Baryon 1980), edited by N. Isgur, Toronto 1980.